

POWERED BY **Dialog****Dialog eLink:** Order File History**IMIDAZOLIDINE DERIVATIVE****Publication Number:** 02-019363 (JP 2019363 A)**Published:** January 23, 1990**Inventors:**

- HENMI KEIJI
- SHIMA ICHIRO
- FUKAMI NAOKI
- HASHIMOTO SHINJI

Applicants

- FUJISAWA PHARMACEUT CO LTD (A Japanese Company or Corporation), JP (Japan)

Application Number: 63-169900 (JP 88169900)**Filed:** July 06, 1988**International Class (IPC Edition 5):**

- C07D-233/72
- C07D-233/74
- C07D-403/06
- A61K-031/415
- A61K-031/415
- A61K-031/415
- A61K-031/415
- A61K-031/415

JAPIO Class:

- 14.1 (ORGANIC CHEMISTRY--- Organic Compounds)
- 14.4 (ORGANIC CHEMISTRY--- Medicine)

Abstract:

NEW MATERIAL: An imidazolidine derivative shown by formula I (R(sup 1) is H or (lower) alkyl; R(sup 2) is lower alkyl, lower alkylthio (lower) alkyl or tri-(lower) alkoxyaryl; R(sup 3) is benzyl; R(sup 4) is lower alkyl or ar-(lower) alkyl; or R(sup 3) is H, phenyl (2C-6C) alkyl, naphthyl (lower) alkyl, aryl, diphenyl (lower) alkyl, etc.; R(sup 4) is H or lower alkyl; with the proviso that when R(sup 3) and R(sup 4) are H, R(sup 1) is lower alkyl) and a salt thereof.

EXAMPLE: 3-Butyl-5-(1-phenylethyl)imidazolidine-2,4-dione.

USE: Useful as an antagonist of platelet activating factor. A preventive and remedy for diseases such as allergic symptoms and thrombosis.

PREPARATION: For example, as shown the reaction formula, a compound shown by formula II is reacted with a

compound shown by formula III to give a compound shown by formula Ia among compounds shown by formula I.
(From: *Patent Abstracts of Japan*, Section: C, Section No. 705, Vol. 14, No. 149, Pg. 56, March 22, 1990)

JAPIO

© 2009 Japan Patent Information Organization. All rights reserved.

Dialog® File Number 347 Accession Number 3043863